

# Memorandum Maths 2014 P1

The Hamiltonian  $H(u, u_x)$  defined on functions on  $\mathbb{R} \times X$ , where  $X$  is a compact manifold, has critical points which are solutions of the linear Klein-Gordon equation. The author considers perturbations of this Hamiltonian, given by polynomial expressions depending on first order derivatives of  $u$ . The associated PDE is then a quasi-linear Klein-Gordon equation. The author shows that, when  $X$  is the sphere, and when the mass parameter  $m$  is outside an exceptional subset of zero measure, smooth Cauchy data of small size give rise to almost global solutions, i.e. solutions defined on a time interval of length  $cN^{-1}$  for any  $N$ . Previous results were limited either to

## Download Ebook Memorandum Maths 2014 P1

the semi-linear case (when the perturbation of the Hamiltonian depends only on  $u$ ) or to the one dimensional problem. The proof is based on a quasi-linear version of the Birkhoff normal forms method, relying on convenient generalizations of para-differential calculus.

Hua's fundamental theorem of geometry of matrices describes the general form of bijective maps on the space of all  $m \times n$  matrices over a division ring  $\mathbb{D}$  which preserve adjacency in both directions. Motivated by several applications the author studies a long standing open problem of possible improvements. There are three natural questions. Can we replace the assumption of preserving adjacency in both directions by the weaker assumption of preserving adjacency in one direction only and still get the

## Download Ebook Memorandum Maths 2014 P1

same conclusion? Can we relax the bijectivity assumption? Can we obtain an analogous result for maps acting between the spaces of rectangular matrices of different sizes? A division ring is said to be EAS if it is not isomorphic to any proper subring. For matrices over EAS division rings the author solves all three problems simultaneously, thus obtaining the optimal version of Hua's theorem. In the case of general division rings he gets such an optimal result only for square matrices and gives examples showing that it cannot be extended to the non-square case.

Let  $\rho$  be the automorphic representation of  $GL_2$  generated by a full level cuspidal Siegel eigenform that is not a Saito-Kurokawa lift, and  $\sigma$  be an arbitrary cuspidal, automorphic representation of  $GL_2$ . Using Furusawa's integral representation for combined

## Download Ebook Memorandum Maths 2014 P1

with a pullback formula involving the unitary group  $U_n$ , the authors prove that the  $L$ -functions are "nice". The converse theorem of Cogdell and Piatetski-Shapiro then implies that such representations have a functorial lifting to a cuspidal representation of  $GL_n$ . Combined with the exterior-square lifting of Kim, this also leads to a functorial lifting of  $L$  to a cuspidal representation of  $GL_n$ . As an application, the authors obtain analytic properties of various  $L$ -functions related to full level Siegel cusp forms. They also obtain special value results for  $L$  and  $\zeta$ .

A stationary solution of the rotating Navier-Stokes equations with a boundary condition is called an Ekman boundary layer. This book constructs stationary solutions of the rotating Navier-Stokes-Boussinesq equations with

## Download Ebook Memorandum Maths 2014 P1

stratification effects in the case when the rotating axis is not necessarily perpendicular to the horizon. The author calls such stationary solutions Ekman layers. This book shows the existence of a weak solution to an Ekman perturbed system, which satisfies the strong energy inequality. Moreover, the author discusses the uniqueness of weak solutions and computes the decay rate of weak solutions with respect to time under some assumptions on the Ekman layers and the physical parameters. The author also shows that there exists a unique global-in-time strong solution of the perturbed system when the initial datum is sufficiently small. Comparing a weak solution satisfying the strong energy inequality with the strong solution implies that the weak solution is smooth with respect to time when time is

sufficiently large.

The Grothendieck Inequality Revisited

Semiclassical Standing Waves with Clustering Peaks for

Nonlinear Schrödinger Equations

A Power Law of Order  $1/4$  for Critical Mean Field Swendsen-

Wang Dynamics

Reflections on Continuing Professional Development of

Mathematics Teachers in a Collaborative Partnership

Oswaal CBSE 5 Years' Solved Papers, Class 10 (English Lang.

& Lit., Hindi-A, Hindi-B, Sanskrit, Social Science, Science

Mathematics (Standard + Basic) (For 2022 Exam)

***The authors develop elements of a general  
dilation theory for operator-valued measures.***

***Hilbert space operator-valued measures are***

***closely related to bounded linear maps on abelian von Neumann algebras, and some of their results include new dilation results for bounded linear maps that are not necessarily completely bounded, and from domain algebras that are not necessarily abelian. In the non-cb case the dilation space often needs to be a Banach space. They give applications to both the discrete and the continuous frame theory. There are natural associations between the theory of frames (including continuous frames and framings), the theory of operator-valued measures on sigma-algebras of sets, and the theory of continuous linear maps between***

***-algebras. In this connection frame theory itself is identified with the special case in which the domain algebra for the maps is an abelian von Neumann algebra and the map is normal (i.e. ultraweakly, or weakly, or  $w^*$ ) continuous.***

***Caught in the Act Reflections on Continuing Professional Development of Mathematics Teachers in a Collaborative Partnership***  
**SUN MeDIA**

***Spectral triples for nonunital algebras model locally compact spaces in noncommutative geometry. In the present text, the authors prove the local index formula for spectral triples over nonunital algebras, without the***



***assumption of local units in our algebra. This formula has been successfully used to calculate index pairings in numerous noncommutative examples. The absence of any other effective method of investigating index problems in geometries that are genuinely noncommutative, particularly in the nonunital situation, was a primary motivation for this study and the authors illustrate this point with two examples in the text. In order to understand what is new in their approach in the commutative setting the authors prove an analogue of the Gromov-Lawson relative index formula (for Dirac type operators) for even dimensional manifolds with***

***bounded geometry, without invoking compact supports. For odd dimensional manifolds their index formula appears to be completely new. The author studies a family of renormalization transformations of generalized diamond hierarchical Potts models through complex dynamical systems. He proves that the Julia set (unstable set) of a renormalization transformation, when it is treated as a complex dynamical system, is the set of complex singularities of the free energy in statistical mechanics. He gives a sufficient and necessary condition for the Julia sets to be disconnected. Furthermore, he proves that all Fatou***

***components (components of the stable sets) of this family of renormalization transformations are Jordan domains with at most one exception which is completely invariant. In view of the problem in physics about the distribution of these complex singularities, the author proves here a new type of distribution: the set of these complex singularities in the real temperature domain could contain an interval. Finally, the author studies the boundary behavior of the first derivative and second derivative of the free energy on the Fatou component containing the infinity. He also gives an explicit value of the second order critical exponent of the free***

***energy for almost every boundary point.***

***A Complete Classification of the Isolated Singularities for Nonlinear Elliptic Equations with Inverse Square Potentials***

***Higher-Order Time Asymptotics of Fast Diffusion in Euclidean Space: A Dynamical Systems Approach***

***Quaternionic Contact Einstein Structures and the Quaternionic Contact Yamabe Problem***

***Index Theory for Locally Compact Noncommutative Geometries***

***New Complex Analytic Methods in the Study of Non-Orientable Minimal Surfaces in  $R^n$***

***Joseph and Hodges–Levasseur (in the A case)***

## Download Ebook Memorandum Maths 2014 P1

*described the spectra of all quantum function algebras on simple algebraic groups in terms of the centers of certain localizations of quotients of by torus invariant prime ideals, or equivalently in terms of orbits of finite groups. These centers were only known up to finite extensions. The author determines the centers explicitly under the general conditions that the deformation parameter is not a root of unity and without any restriction on the characteristic of the ground field. From it he deduces a more explicit description of all prime ideals of than the previously known ones and an*

## Download Ebook Memorandum Maths 2014 P1

*explicit parametrization of .*

*• Solved Board Examination Paper 2020 along with CBSE Marking Scheme from 2016 to 2019 for in-depth study. • Previous Years' Board Examination Questions with Solutions from March 2016 to March 2019 to facilitate focused study. • Handwritten Toppers' Answer sheets from 2016-2019 for perfection in answering Board Examination Questions • Detailed answers have been provided wherever necessary for clarity of concepts • Hybrid edition for Digital Learning*

*The authors study the following singularly perturbed problem: in . Their main result is*

## Download Ebook Memorandum Maths 2014 P1

*the existence of a family of solutions with peaks that cluster near a local maximum of . A local variational and deformation argument in an infinite dimensional space is developed to establish the existence of such a family for a general class of nonlinearities . Consider a Hamiltonian action of a compact connected Lie group on a symplectic manifold . Conjecturally, under suitable assumptions there exists a morphism of cohomological field theories from the equivariant Gromov-Witten theory of to the Gromov-Witten theory of the symplectic quotient. The morphism should be a deformation of the Kirwan map.*

## Download Ebook Memorandum Maths 2014 P1

*The idea, due to D. A. Salamon, is to define such a deformation by counting gauge equivalence classes of symplectic vortices over the complex plane . The present memoir is part of a project whose goal is to make this definition rigorous. Its main results deal with the symplectically aspherical case.*

*Cohomology for Quantum Groups via the Geometry of the Nullcone*

*New Foundations for Geometry: Two Non-Additive Languages for Arithmetical Geometry*

*Nonlinear Stability of Ekman Boundary Layers in Rotating Stratified Fluids*

*Special Values of Automorphic Cohomology*



## **Classes**

### **Generalized Descriptive Set Theory and Classification Theory**

*The research reported in Caught in the Act draws together the insights gained from a continuous professional development initiative for high school mathematics teachers, teaching in low socio-economic status and socially challenged environments in South Africa. The chapters show the possibilities for enhancing achievement in school mathematics if a strength-based approach is adopted to develop teaching with teachers and when their concerns are taken seriously. The book demonstrates that the ecological*

*relevance – fitness for the context in which teachers are teaching – of the “goods” developed and distributed by initiatives should be a major consideration to offer hope for sustainable implementation to improve school mathematics education. The book is of interest to mathematics teachers, school leaders, mathematics curriculum advisors, policy makers and researchers.*

*Descriptive set theory is mainly concerned with studying subsets of the space of all countable binary sequences. In this paper the authors study the generalization where countable is replaced by uncountable. They explore properties of generalized Baire and Cantor spaces,*

*equivalence relations and their Borel reducibility. The study shows that the descriptive set theory looks very different in this generalized setting compared to the classical, countable case. They also draw the connection between the stability theoretic complexity of first-order theories and the descriptive set theoretic complexity of their isomorphism relations. The authors' results suggest that Borel reducibility on uncountable structures is a model theoretically natural way to compare the complexity of isomorphism relations. The classical Grothendieck inequality is viewed as a statement about representations of functions of two variables over discrete domains by integrals of two-fold*

*products of functions of one variable. An analogous statement is proved, concerning continuous functions of two variables over general topological domains. The main result is the construction of a continuous map  $\Phi$  from  $L^2(A)$  into  $L^2(\Omega_A, \mathbb{P}_A)$ , where  $A$  is a set,  $\Omega_A = \{-1, 1\}^A$ , and  $\mathbb{P}_A$  is the uniform probability measure on  $\Omega_A$ .*

*In this monograph the authors introduce a new method to study bifurcations of KAM tori with fixed Diophantine frequency in parameter-dependent Hamiltonian systems. It is based on Singularity Theory of critical points of a real-*

*valued function which the authors call the potential. The potential is constructed in such a way that: nondegenerate critical points of the potential correspond to twist invariant tori (i.e. with nondegenerate torsion) and degenerate critical points of the potential correspond to non-twist invariant tori. Hence, bifurcating points correspond to non-twist tori.*

*A Quantum Kirwan Map: Bubbling and Fredholm Theory for Symplectic Vortices over the Plane*

*Spectra of Symmetrized Shuffling Operators*

*Critical Population and Error Threshold on the Sharp Peak Landscape for a Moran Model*

*Shock Waves in Conservation Laws with Physical Viscosity*

### *The Optimal Version of Hua's Fundamental Theorem of Geometry of Rectangular Matrices*

The author develops a homology theory for Smale spaces, which include the basics sets for an Axiom A diffeomorphism. It is based on two ingredients. The first is an improved version of Bowen's result that every such system is the image of a shift of finite type under a finite-to-one factor map. The second is Krieger's dimension group invariant for shifts of

finite type. He proves a Lefschetz formula which relates the number of periodic points of the system for a given period to trace data from the action of the dynamics on the homology groups. The existence of such a theory was proposed by Bowen in the 1970s. The authors investigate the global continuity on spaces with of Fourier integral operators with smooth and rough amplitudes and/or phase functions subject to certain necessary non-

degeneracy conditions. In this context they prove the optimal global boundedness result for Fourier integral operators with non-degenerate phase functions and the most general smooth Hörmander class amplitudes i.e. those in with . They also prove the very first results concerning the continuity of smooth and rough Fourier integral operators on weighted spaces, with and (i.e. the Muckenhoupt weights) for operators with rough and smooth



amplitudes and phase functions satisfying a suitable rank condition. The authors develop a theory for the existence of perfect matchings in hypergraphs under quite general conditions. Informally speaking, the obstructions to perfect matchings are geometric, and are of two distinct types: 'space barriers' from convex geometry, and 'divisibility barriers' from arithmetic lattice-based constructions. To formulate precise

results, they introduce the setting of simplicial complexes with minimum degree sequences, which is a generalisation of the usual minimum degree condition. They determine the essentially best possible minimum degree sequence for finding an almost perfect matching. Furthermore, their main result establishes the stability property: under the same degree assumption, if there is no perfect matching then there must be a space or

divisibility barrier. This allows the use of the stability method in proving exact results. Besides recovering previous results, the authors apply our theory to the solution of two open problems on hypergraph packings: the minimum degree threshold for packing tetrahedra in  $k$ -graphs, and Fischer's conjecture on a multipartite form of the Hajnal-Szemerédi Theorem. Here they prove the exact result for tetrahedra and the asymptotic result for Fischer's

conjecture; since the exact result for the latter is technical they defer it to a subsequent paper.

The authors study the complex geometry and coherent cohomology of nonclassical Mumford-Tate domains and their quotients by discrete groups. Their focus throughout is on the domains which occur as open  $\mathbb{C}^*$ -orbits in the flag varieties for  $SO^*(2n)$  and  $SO^*(2n+1)$ , regarded as classifying spaces for Hodge structures of weight three. In the context

provided by these basic examples, the authors formulate and illustrate the general method by which correspondence spaces give rise to Penrose transforms between the cohomologies of distinct such orbits with coefficients in homogeneous line bundles.

Formality of the Little  $N$ -disks

Operad

Taxonomy of Educational Objectives

Transfer of Siegel Cusp Forms of Degree

2

Near Soliton Evolution for Equivariant Schrödinger Maps in Two Spatial Dimensions Ioan Bejenaru, University of California, San Diego, La Jolla, CA, and Daniel Tataru, University of California, Berkeley, Berkeley, CA  
Local Entropy Theory of a Random Dynamical System

***The authors consider the Schrödinger Map equation in 2+1 dimensions, with values into  $\mathbb{S}^2$ . This admits a lowest energy steady state  $Q$ , namely the stereographic projection, which extends to a two***

**dimensional family of steady states by scaling and rotation. The authors prove that  $Q$  is unstable in the energy space  $\dot{H}^1$ . However, in the process of proving this they also show that within the equivariant class  $Q$  is stable in a stronger topology  $X \subset \dot{H}^1$ .**

**For a finite real reflection group  $W$  and a  $W$ -orbit  $\mathcal{O}$  of flats in its reflection arrangement--or equivalently a conjugacy class of its parabolic subgroups--the authors introduce a statistic  $\text{noninv}_{\mathcal{O}}(w)$  on  $w$  in  $W$  that counts the number of " $\mathcal{O}$ -noninversions" of  $w$ . This generalizes the classical (non-)inversion statistic for**

**permutations  $w$  in the symmetric group  $\frac{S}{n}$ . The authors then study the operator  $\nu_{\mathcal{O}}$  of right-multiplication within the group algebra  $\mathbb{C} W$  by the element that has**

**$\operatorname{noninv}_{\mathcal{O}}(w)$  as its coefficient on  $w$ .**

**The authors study the perturbation of a shock wave in conservation laws with physical viscosity. They obtain the detailed pointwise estimates of the solutions. In particular, they show that the solution converges to a translated shock profile. The strength of the perturbation and that of the shock are assumed to be small but independent. The authors'**



***assumptions on the viscosity matrix are general so that their results apply to the Navier-Stokes equations for the compressible fluid and the full system of magnetohydrodynamics, including the cases of multiple eigenvalues in the transversal fields, as long as the shock is classical. The authors' analysis depends on accurate construction of an approximate Green's function. The form of the ansatz for the perturbation is carefully constructed and is sufficiently tight so that the author can close the nonlinear term through Duhamel's principle. Let  $F$  be a non-Archimedean local field. Let  $\mathcal{W}_F$  be the Weil group of  $F$  and  $\mathcal{P}_F$  the wild inertia subgroup of***

$\mathcal{W}_F$ . Let  $\widehat{\mathcal{W}}_F$  be the set of equivalence classes of irreducible smooth representations of  $\mathcal{W}_F$ . Let  $\mathcal{A}^0_n(F)$  denote the set of equivalence classes of irreducible cuspidal representations of  $\mathrm{GL}_n(F)$  and set  $\widehat{\mathrm{GL}}_F = \bigcup_{n \geq 1} \mathcal{A}^0_n(F)$ . If  $\sigma \in \widehat{\mathcal{W}}_F$ , let  $^L\sigma \in \widehat{\mathrm{GL}}_F$  be the cuspidal representation matched with  $\sigma$  by the Langlands Correspondence. If  $\sigma$  is totally wildly ramified, in that its restriction to  $\mathcal{P}_F$  is irreducible, the authors treat  $^L\sigma$  as

**known. From that starting point, the authors construct an explicit bijection  $\mathbb{N}:\widehat{\mathcal{W}}_F$  to  $\widehat{\mathrm{GL}}_F$ , sending  $\sigma$  to  $^N\sigma$ . The authors compare this "naïve correspondence" with the Langlands correspondence and so achieve an effective description of the latter, modulo the totally wildly ramified case. A key tool is a novel operation of "internal twisting" of a suitable representation  $\pi$  (of  $\mathcal{W}_F$  or  $\mathrm{GL}_n(F)$ ) by tame characters of a tamely ramified field extension of  $F$ , canonically associated to  $\pi$ . The authors show this operation is preserved by the Langlands correspondence.**

***The Classification of Educational Goals***

***A Geometric Theory for Hypergraph Matching***

***Global and Local Regularity of Fourier Integral***

***Operators on Weighted and Unweighted Spaces***

***Effective Hamiltonians for Constrained Quantum***

***Systems***

***Operator-Valued Measures, Dilations, and the Theory of Frames***

*The authors define combinatorial Floer homology of a transverse pair of noncontractible nonisotopic embedded loops in an oriented  $3$ -manifold without boundary, prove that it is invariant under isotopy, and prove that it is isomorphic to the original*

## Download Ebook Memorandum Maths 2014 P1

*Lagrangian Floer homology. Their proof uses a formula for the Viterbo–Maslov index for a smooth lune in a  $-$ -manifold.*

*The authors consider the time-dependent Schrödinger equation on a Riemannian manifold with a potential that localizes a certain subspace of states close to a fixed submanifold . When the authors scale the potential in the directions normal to by a parameter , the solutions concentrate in an  $-$ -neighborhood of . This situation occurs for example in quantum wave guides and for the motion of nuclei in electronic potential surfaces in quantum molecular dynamics. The*

authors derive an effective Schrödinger equation on the submanifold and show that its solutions, suitably lifted to  $M$ , approximate the solutions of the original equation on  $M$  up to errors of order  $\epsilon$  at time  $t$ . Furthermore, the authors prove that the eigenvalues of the corresponding effective Hamiltonian below a certain energy coincide up to errors of order  $\epsilon$  with those of the full Hamiltonian under reasonable conditions.

All the new tools mentioned above apply to non-orientable minimal surfaces endowed with a fixed choice of a conformal structure. This enables the authors to obtain significant new

*applications to the global theory of non-orientable minimal surfaces. In particular, they construct proper non-orientable conformal minimal surfaces in  $R^n$  with any given conformal structure, complete non-orientable minimal surfaces in  $R^n$  with arbitrary conformal type whose generalized Gauss map is nondegenerate and omits  $n$  hyperplanes of  $CP^{n-1}$  in general position, complete non-orientable minimal surfaces bounded by Jordan curves, and complete proper non-orientable minimal surfaces normalized by bordered surfaces in  $p$ -convex domains of  $R^n$ . The little -disks operad, , along with its*

## Download Ebook Memorandum Maths 2014 P1

variants, is an important tool in homotopy theory. It is defined in terms of configurations of disjoint  $n$ -dimensional disks inside the standard unit disk in  $\mathbb{R}^n$  and it was initially conceived for detecting and understanding  $n$ -fold loop spaces. Its many uses now stretch across a variety of disciplines including topology, algebra, and mathematical physics. In this paper, the authors develop the details of Kontsevich's proof of the formality of little  $n$ -disks operad over the field of real numbers. More precisely, one can consider the singular chains on  $\mathcal{L}_n$  as well as the singular homology of



. These two objects are operads in the category of chain complexes. The formality then states that there is a zig-zag of quasi-isomorphisms connecting these two operads. The formality also in some sense holds in the category of commutative differential graded algebras. The authors additionally prove a relative version of the formality for the inclusion of the little  $n$ -disks operad in the little  $m$ -disks operad when  $n \leq m$ .

*Quasi-Linear Perturbations of Hamiltonian Klein-Gordon Equations on Spheres*

*Combinatorial Floer Homology*

*To an Effective Local Langlands*

## *Correspondence*

### *A Homology Theory for Smale Spaces*

### *Weighted Bergman Spaces Induced by Rapidly Increasing Weights*

This paper quantifies the speed of convergence and higher-order asymptotics of fast diffusion dynamics on  $\mathbb{R}^n$  to the Barenblatt (self similar) solution.

Degeneracies in the parabolicity of this equation are cured by re-expressing the dynamics on a manifold with a cylindrical end, called the cigar. The nonlinear evolution becomes differentiable in Hölder

## Download Ebook Memorandum Maths 2014 P1

spaces on the cigar. The linearization of the dynamics is given by the Laplace-Beltrami operator plus a transport term (which can be suppressed by introducing appropriate weights into the function space norm), plus a finite-depth potential well with a universal profile. In the limiting case of the (linear) heat equation, the depth diverges, the number of eigenstates increases without bound, and the continuous spectrum recedes to infinity. The authors provide a detailed study of the linear and nonlinear problems

## Download Ebook Memorandum Maths 2014 P1

in Hölder spaces on the cigar, including a sharp boundedness estimate for the semigroup, and use this as a tool to obtain sharp convergence results toward the Barenblatt solution, and higher order asymptotics. In finer convergence results (after modding out symmetries of the problem), a subtle interplay between convergence rates and tail behavior is revealed. The difficulties involved in choosing the right functional spaces in which to carry out the analysis can be interpreted as genuine features of the

## Download Ebook Memorandum Maths 2014 P1

equation rather than mere annoying technicalities.

Polynomial approximation on convex polytopes in  $\mathbb{R}^n$  is considered in uniform and  $L^p$ -norms. For an appropriate modulus of smoothness matching direct and converse estimates are proven. In the  $L^p$ -case so called strong direct and converse results are also verified. The equivalence of the moduli of smoothness with an appropriate  $\Psi$ -functional follows as a consequence. The results solve a problem that was left open since the mid 1980s when some of the

## Download Ebook Memorandum Maths 2014 P1

present findings were established for special, so-called simple polytopes. The goal of this work is to propose a finite population counterpart to Eigen's model, which incorporates stochastic effects. The author considers a Moran model describing the evolution of a population of size of chromosomes of length over an alphabet of cardinality  $k$ . The mutation probability per locus is  $\mu$ . He deals only with the sharp peak landscape: the replication rate is  $1$  for the master sequence and  $1 - \mu$  for the other sequences. He

## Download Ebook Memorandum Maths 2014 P1

studies the equilibrium distribution of  
the process in the regime where  
Introduction Statement of the results  
Mixing time preliminaries Outline of the  
proof of Theorem 2.1 Random graph  
estimates Supercritical case Subcritical  
case Critical Case Fast mixing of the  
Swendsen-Wang process on trees  
Acknowledgements Bibliography  
Relative Equilibria in the 3-Dimensional  
Curved n-Body Problem

Polynomial Approximation on Polytopes

## Download Ebook Memorandum Maths 2014 P1

Julia Sets and Complex Singularities of  
Free Energies

Singularity Theory for Non-Twist KAM Tori

To view the abstract go to

<http://www.ams.org/books/memo/1166>.

A partial solution of the quaternionic contact Yamabe problem on the quaternionic sphere is given. It is shown that the torsion of the Biquard connection vanishes exactly when the trace-free part of the horizontal Ricci tensor of the Biquard connection is zero and this occurs precisely on 3-Sasakian manifolds. All conformal transformations sending the standard flat torsion-free quaternionic contact



structure on the quaternionic Heisenberg group to a quaternionic contact structure with vanishing torsion of the Biquard connection are explicitly described. A "3-Hamiltonian form" of infinitesimal conformal automorphisms of quaternionic contact structures is presented.

In this paper the authors extend the notion of a continuous bundle random dynamical system to the setting where the action of  $\mathbb{R}$  or  $\mathbb{N}$  is replaced by the action of an infinite countable discrete amenable group. Given such a system, and a monotone sub-additive invariant family of random continuous functions, they introduce the concept of local

fiber topological pressure and establish an associated variational principle, relating it to measure-theoretic entropy. They also discuss some variants of this variational principle. The authors introduce both topological and measure-theoretic entropy tuples for continuous bundle random dynamical systems, and apply variational principles to obtain a relationship between these of entropy tuples. Finally, they give applications of these results to general topological dynamical systems, recovering and extending many recent results in local entropy theory.

Operator Theory, Operator Algebras, and Applications  
Caught in the Act

## On the Spectra of Quantum Groups